

Management Plan for the Western Toad (*Anaxyrus boreas*), Calling and Non-calling populations, in Canada

Western Toad



2020



Government
of Canada

Gouvernement
du Canada

Canada

Recommended citation:

Environment and Climate Change Canada. 2020. Management Plan for the Western Toad (*Anaxyrus boreas*), Calling and Non-calling populations, in Canada¹. *Species at Risk Act* Management Plan Series. Environment and Climate Change Canada, Ottawa. v + 39 pp.

Official version

The official version of the recovery documents is the one published in PDF. All hyperlinks were valid as of date of publication.

Non-official version

The non-official version of the recovery documents is published in HTML format and all hyperlinks were valid as of date of publication.

For copies of the management plan, or for additional information on species at risk, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Status Reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk \(SAR\) Public Registry](#)².

Cover illustration: copyright Syd Cannings

Également disponible en français sous le titre
« Plan de gestion du crapaud de l'Ouest (*Anaxyrus boreas*), population chantante et population non-chantante, au Canada »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Environment and Climate Change, 2020. All rights reserved.

ISBN 978-0-660-35729-4

Catalogue no. En3-5/107-2020E-PDF

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

¹ At the time of proposed posting, in 2016, the document title did not include the two population names and was entitled *Management Plan for the Western Toad (Anaxyrus boreas), in Canada*. In 2018, Schedule 1 of SARA was amended to include both populations of this species and the title of this final document has been changed accordingly.

² www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html

Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#)³ agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29)(SARA), the federal competent ministers are responsible for the preparation of management plans for listed species of special concern and are required to report on progress within five years after the publication of the final document on the Species at Risk (SAR) Public Registry.

The Minister of Environment and Climate Change and Minister responsible for the Parks Canada Agency is the competent minister under SARA for the Western Toad, Calling and Non-calling populations, and has prepared this management plan as per section 65 of SARA. To the extent possible, it has been prepared in cooperation with the provinces of British Columbia and Alberta, and Yukon and Northwest Territories as per section 66(1) of SARA.

Success in the conservation of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this plan and will not be achieved by Environment and Climate Change Canada, the Parks Canada Agency, or any other jurisdiction alone. All Canadians are invited to join in supporting and implementing this plan for the benefit of the Western Toad, Calling and Non-calling populations, and Canadian society as a whole.

Implementation of this management plan is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

³ www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2

Acknowledgments

An earlier draft of this document was prepared by Kristiina Ovaska and Lennart Sopuck (Biolinx Environmental Research Ltd.), with input from representatives of federal, provincial, and territorial governments, and regional experts. The document was updated by Greg Ferguson, Holly Middleton, Megan Harrison, Eric Gross, and Jared Maida of Environment and Climate Change Canada, with review and input from the following people: Lucy Reiss, Syd Cannings, David Cunnington, Andy Didiuk, Mike Gill, Stephen Hureau, Donna Mulders, Mark Wayland, and Donna Bigelow (Environment and Climate Change Canada, Canadian Wildlife Service); Diane Casimir, Barb Johnston, Rhonda Kindopp, Lisa Larson, Charlie Pacas, Derek Petersen, Brenda Shepherd, Doug Tate, and Carmen Wong (Parks Canada); Tom Jung, Tess McLeod, Bruce Bennett and Todd Powell (Government of Yukon); Rob Gau, Nic Larter, and Joanna Wilson (Government of Northwest Territories); Purnima Govindarajulu, James Quayle, Leah Westereng, Doug Wilson, Connie Miller Retzer, Sarma Liepins, Dave Trotter, and Peter Fielder (Government of BC); Gavin Berg, Lisa Wilkinson, and Scott Stevens (Government of Alberta); Brian Eaton (Alberta Innovates - Technology Futures, Vegreville); Cindy Paszkowski (University of Alberta); Danna Schock (Detroit Zoological Society), Brian Slough (Whitehorse, YT); Doug Whiteside (Calgary Zoo); and Elke Wind (Nanaimo, BC).

Executive Summary

The Western Toad (*Anaxyrus boreas*) is a large, stocky toad. It ranges in colour from greenish to tan, brown, or black with a light line down the centre of its back and a pronounced paratoid (cheek) gland. The Western Toad was assessed as Special Concern by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2002 and listed under Schedule 1 of the *Species at Risk Act* in 2005. COSEWIC reconfirmed its status in 2012 and divided the species into two populations or designatable units: Non-calling and Calling. In 2018, Schedule 1 of SARA was amended to reflect this same division.

The Western Toad has a wide distribution in western North America, from Baja California to Alaska, and from the Pacific Coast to Colorado and Alberta. In Canada, the Western Toad occurs throughout the majority of British Columbia, western Alberta, and southeastern Yukon and southwestern Northwest Territories. The majority of the Non-calling population in Canada occurs within British Columbia and almost the entire global range of the Calling population is within Alberta.

The Non-calling population has suffered declines and extirpations in the United States and declines appear to have occurred in Canada along the south coast of British Columbia and in other localized areas within the province over a number of decades. In Yukon and Northwest Territories, the number of known occurrences has increased (likely due to increased survey effort), but there has been no evidence of expansion outside the historical range. The Calling population remains widespread throughout much of its historical range in Alberta and may be expanding its range eastward. However, declines are suspected and also projected based on known vulnerabilities and threats.

The Western Toad uses a wide variety of aquatic habitats for breeding and terrestrial habitats for foraging and hibernation. These habitats may be several kilometers apart, which requires Western Toads to move extensively and increases their vulnerability to human developments and activities. High breeding site fidelity, communal hibernation and egg-laying, and the tendency for newly emerged Western Toads to form large post-metamorphic aggregations also increase their vulnerability. The species' reliance on high adult survival to sustain populations through periods of poor reproductive success means that threats that impact adult survival can have particularly pronounced effects.

The main threats to both populations are transportation and service corridors (habitat loss/fragmentation and road mortality), and invasive and other problematic species and pathogens (particularly, infection with the amphibian chytrid fungus [*Batrachochytrium dendrobatidis*]). Other threats common to both populations include logging/wood harvesting, pollution, and climate change. The Calling population also is threatened by agricultural activities and oil and gas drilling.

The management objective is to maintain stable or increasing populations distributed throughout the species' present range in Canada.

The broad strategies and conservation measures to achieve the management objective are outlined in Section 6.2 and 6.3 of this document.

Table of Contents

Preface.....	I
Acknowledgments	II
Executive Summary	III
1. COSEWIC Species Assessment Information.....	1
2. Species Status Information	2
3. Species Information	3
3.1. Species Description.....	3
3.2. Population and Distribution.....	6
3.3. Needs of the Western Toad.....	10
4. Threats.....	12
4.1. Threat Assessment.....	13
4.2. Description of Threats	16
5. Management Objective	20
6. Broad Strategies and Conservation Measures	20
6.1 Actions Already Completed or Currently Underway.....	20
6.2 Broad Strategies.....	24
6.3 Conservation Measures.....	25
7. Measuring Progress.....	26
8. References	27
9. Personal Communications	37
Appendix A: Effects on the Environment and Other Species	39

1. COSEWIC* Species Assessment Information

Date of Assessment: November 2012

Common Name (population): Western Toad - Non-calling population

Scientific Name: *Anaxyrus boreas*

COSEWIC Status: Special Concern

Reason for Designation: This species has suffered population declines and population extirpations in the southern part of its range in British Columbia, as well as in the USA. The toads are particularly sensitive to emerging skin disease caused by the amphibian chytrid fungus, which has been linked to global amphibian declines. It is relatively intolerant of urban expansion, conversion of habitat for agricultural use, and habitat fragmentation resulting from resource extraction and road networks. Life history characteristics, including infrequent breeding by females, aggregation at communal, traditionally used breeding sites, and migrations to and from breeding sites, make populations vulnerable to habitat degradation and fragmentation. The species remains widespread, but declines are suspected and projected based on known vulnerabilities and threats.

Canadian Occurrence: Yukon, Northwest Territories, British Columbia, Alberta

COSEWIC Status History: The species was considered a single unit and designated Special Concern in November 2002. Split into two populations in November 2012. The Non-calling population was designated Special Concern in November 2012.

* COSEWIC (Committee on the Status of Endangered Wildlife in Canada)

Date of Assessment: November 2012

Common Name (population): Western Toad - Calling population

Scientific Name: *Anaxyrus boreas*

COSEWIC Status: Special Concern

Reason for Designation: Almost the entire range of the calling population is within Canada. The toads are particularly sensitive to emerging skin disease caused by the amphibian chytrid fungus, which has been linked to global amphibian declines. This species is relatively intolerant of urban expansion, conversion of habitat for agricultural use, and habitat fragmentation resulting from resource extraction and road networks. Life history characteristics, including infrequent breeding by females, aggregation at communal, traditionally used breeding sites, and migrations to and from breeding sites, make populations vulnerable to habitat degradation and fragmentation. The species remains widespread throughout much of their historic range in Alberta and may be expanding their range eastwards. However, declines are suspected and projected based on known vulnerabilities and threats.

Canadian Occurrence: British Columbia, Alberta

COSEWIC Status History: The species was considered a single unit and designated Special Concern in November 2002. Split into two populations in November 2012. The Calling population was designated Special Concern in November 2012.

2. Species Status Information

Status ranks for the Western Toad are listed in Table 1. The International Union for Conservation of Nature (IUCN) has designated the species as “near threatened” due to declines and extirpations over parts of its distribution in the United States (IUCN 2014). COSEWIC assessed the Western Toad as Special Concern (COSEWIC 2002) and it was listed under Schedule 1 of the Canadian *Species at Risk Act* (SARA) in 2005. COSEWIC reconfirmed its status in 2012 and divided the species into two populations or designatable units: Non-calling and Calling (COSEWIC 2012). In 2018, Schedule 1 of SARA was amended to include this same division. Approximately 40% of the species’ distribution is within Canada. In the Northwest Territories, the Western Toad is listed as a threatened species (Species at Risk Committee 2014).

Table 1. List and description of various conservation status ranks for Western Toad (British Columbia Ministry of Environment 2010; NatureServe 2014; British Columbia Conservation Data Centre 2014; Working Group on General Status of NWT Species 2016).

Global (G) Rank*	National (N) Rank*	Sub-national (S) Rank*	COSEWIC Status	B.C. Status Rank	B.C. Conservation Framework***
G4	N4	British Columbia: S3S4 Alberta: S3 Yukon: S3 Northwest Territories: S1S3 Alaska S3S4 California: SNR Colorado: S1 Idaho: S3 Montana: S2 Nevada: S4 New Mexico: S1 Oregon: S3 Utah: S3 Washington: S3 Wyoming: S1	SC (Special Concern)	Blue**	Priority 3 under Goal 1, Priority 2 under Goal 2, and Priority 4 under Goal 3

* Rank 1– critically imperiled; 2– imperiled; 3– vulnerable to extirpation or extinction; 4– apparently secure; 5– secure; SNR: status not reported** Includes any indigenous species or subspecies considered to be of Special Concern (formerly Vulnerable) in British Columbia. Taxa of Special Concern have characteristics that make them particularly sensitive or vulnerable to human activities or natural events. Blue-listed taxa are at risk, but are not Extirpated, Endangered, or Threatened.

*** The three goals of the B.C. Conservation Framework are: 1. Contribute to global efforts for species and ecosystem conservation; 2. Prevent species and ecosystems from becoming at risk; 3. Maintain the diversity of native species and ecosystems

3. Species Information

3.1. Species Description

The Western Toad is a member of the large cosmopolitan family Bufonidae, or true toads. Much of the literature on this species refers to it as *Bufo boreas*, but when the genus *Bufo* was split (Frost et al. 2006), the species was placed into the genus *Anaxyrus*. In Canada, COSEWIC (2012) classified the Western Toad into two populations or designatable units; Non-calling and Calling (Figure 1). These two populations are distinguished by the presence (Calling) or absence (Non-calling) of a vocal sac in males and the production of a true advertisement call, which is characterized by long, high-amplitude trills (Pauly 2008). Vocalizations also are made in the Non-calling population, but they are not equivalent to the high-amplitude advertisement call that distinguishes the Calling population (Pauly 2008).

The Western Toad is a robust toad with an adult body length of 55 - 145 mm (Corkran and Thoms 2006). Its colour ranges from greenish to tan, brown, grey, or black, with or

without mottling; a light line down the centre of the back is usually present, but may be lacking in small toadlets. There is a pronounced oval poison gland (parotoid gland) on each cheek behind the eye and raised poison glands (“warts”) on the back. The Western Toad occurs sympatrically with the superficially similar Canadian Toad (*Anaxyrus hemiophrys*) in eastern Alberta. The Canadian Toad is smaller (adult body length usually 70 mm or less) and has a hump (or boss) on the head between the eyes. Western Toad tadpoles are black and small, roughly 25 - 30 mm total length prior to metamorphosis (Green and Campbell 1984; Blaustein et al. 1995). For illustrations and detailed descriptions of different life stages of the Western Toad see Russell and Bauer 2000; Jones et al. 2005; Corkran and Thoms 2006; and Matsuda et al. 2006.




Figure 1. Sites verified to contain either Non-calling or Calling males of the Western Toad in North America (Pauly 2008 in COSEWIC 2012). Triangles indicate males lacking vocal sacs (white), males with vocal sacs (black), or males with and without vocal sacs (gray). Overall sample size = 1279 individuals (museum and purpose-collected specimens). Numbered sites in Alberta are localities where males were recorded during field visits. The asterisk in northern Alberta denotes individuals observed to have vocal sacs and produce long, pulsed calls, but no specimens or recordings were taken. The grey shaded area is an outdated representation of the global range. A more accurate delineation of the species' range is shown in Figure 2.

3.2. Population and Distribution

3.2.1. Species Distribution

The Western Toad has a wide distribution in western North America, from Baja California to Alaska, and from the Pacific Coast to Colorado and Alberta (Figure 2). Across its range, it occurs from near sea level to elevations of at least 3355 m (NatureServe 2014). In Canada, the Western Toad occurs in British Columbia, Alberta, Yukon, and Northwest Territories (Figure 2). Most of the Canadian distribution is in British Columbia (~70%) and Alberta (~20%), with small portions in the Yukon and Northwest Territories (Figure 2). The Calling population occurs mainly to the east and the Non-calling population to the west of the Rocky Mountains (Pauly 2008; Figure 3), but the boundary is not completely understood and requires additional study (COSEWIC 2012).

In British Columbia, the Western Toad ranges throughout most of the province, including Vancouver Island and the Haida Gwaii archipelago (Matsuda et al. 2006). The species appears to be absent from the Teslin River basin (Government of Yukon 2013) and is likely absent from extreme northeast British Columbia (COSEWIC 2012).

The range of the Western Toad extends slightly into Yukon and Northwest Territories (COSEWIC 2012). In southeast Yukon, it occurs in the southern Liard River basin, where it has been found in five geographically separated localities (Slough and Mennell 2006; Slough 2009a; Yukon CDC 2015). In southwest Northwest Territories, it occurs in the Liard River basin, in the Dehcho region, where it has been found in six different sites (Schock et al. 2009; Species at Risk Committee 2014; Government of the Northwest Territories 2018).

In Alberta, the Western Toad ranges from the forested regions of the southwest to central and northern Alberta, and to a lesser extent into the short-grass prairie and aspen parkland (COSEWIC 2012). The distribution in northern Alberta may be more extensive than illustrated in Figure 3, which reflects limited survey effort, and few historical occurrence records (Russell and Bauer 2000; COSEWIC 2012). Surveys associated with oil and gas development and forestry activities in northern Alberta recently detected Western Toad in new areas, which could be indicative of an eastward range expansion, or simply a reflection of increased survey effort (COSEWIC 2012). If this indicates a true range expansion, the Western Toad may be replacing the Canadian Toad (*Anaxyrus hemiophrys*), which has declined throughout its range in Alberta (COSEWIC 2012).




Figure 2. Global distribution of the Western Toad (dark green shading). Map prepared by Joanna Wilson, Northwest Territories Department of Environment and Natural Resources, 2014; United States and Mexican range based on a map compiled by IUCN, Conservation International, NatureServe, and collaborators in 2004 (NatureServe 2014).




Figure 3. Canadian distribution of the Western Toad (after COSEWIC 2012). The current estimated boundary between the two populations/designatable units (DUs) is indicated by the red line. The Non-calling population is mainly to the west and north of the line and the Calling population is primarily to the east and south.

3.2.2. Population Abundance and Trends

Precise global population and trend data are unavailable, but NatureServe (2014) estimates that the Western Toad population includes at least 100,000 adults (possibly up to 1,000,000) and has declined by up to 50% in the last 200 years (with 10-30% in the past ~18 years [i.e., 3 generations]). In the United States, the Western Toad has suffered widespread declines and disappearances (COSEWIC 2002; NatureServe 2014), partially linked to the epidemic chytridiomycosis, which is caused by the pathogenic chytrid fungus *Batrachochytrium dendrobatidis* (Carey 1993; Muths et al. 2003). Populations in the southern Rocky Mountains and Sierra Nevada have experienced the most severe declines (NatureServe 2014).

Opportunistic observations and localized studies suggest that the Western Toad is widespread, abundant, and persistent across a large portion of its Canadian range. However, scant information exists on population sizes or densities, and few populations have been systematically monitored to evaluate trends (COSEWIC 2012). Congregations at breeding sites and mass migrations of toadlets (metamorphs) may give an appearance of abundance; however, tadpole and metamorph abundance is often a poor indicator of adult population size because breeding success can fluctuate dramatically and juvenile mortality is high (COSEWIC 2012). The Western Toad appears to be less abundant north of 58°N where aggregations of tadpoles and metamorphs have been reported only in the hundreds or thousands (Slough 2004; 2005; 2009a; Schock 2009) compared to elsewhere where aggregations of tens to hundreds of thousands have been reported (COSEWIC 2012).

In British Columbia, numbers appear to be declining along the south mainland coast, Vancouver Island, and in other localized areas (COSEWIC 2012), but rates of decline have not been quantified (Provincial Western Toad Working Group 2014). COSEWIC (2002) reported population declines in the Lower Fraser Valley associated with local anthropogenic habitat loss, and Davis and Gregory (2003) documented the toad's disappearance from a large wetland complex on Vancouver Island. Ohanjanian et al. (2006) visited 87 wetlands in the East Kootenays historically known to support Western Toads and found breeding evidence at less than a third of the sites. In 2011, extant breeding populations were reported in the south coast of the mainland and Vancouver Island (Beasley 2011; Tayless 2011; COSEWIC 2012) although the number of breeding sites is thought to have declined in the past 20 years. Wetland loss has been significant in parts of British Columbia, particularly in the Fraser and Columbia river drainages; in the South Okanagan region (part of the Columbia River drainage), ~85% of the original wetland area has been lost (Austin et al. 2008).

In Alberta, since the 1980s, local residents (J. Russell, pers. comm. 2013) and biologists (Wallis et al. 2002) have observed a declining trend in the numbers of toads in the Waterton Lakes National Park region. Occupancy surveys of breeding sites in Waterton Lakes National Park between 1998 and 2015 showed yearly fluctuations with an increase in occupancy in foothills parkland and montane ecoregions since 2014 (Johnston and Price 2015).. Data collected during pitfall trapping at six monitoring sites

in Alberta from 1997-2008 indicated that Western Toads experienced wide fluctuations in relative abundance and adult abundance trends varied by site: two increasing, one decreasing, and three stable (Eaton and Hiltz 2012). Other reports (cited by COSEWIC (2012), i.e., Eaves 2004; Schank 2008) have indicated that Western Toad populations are either stable or increasing in parts of Alberta. It should be noted that declines or disappearances could have gone unnoticed in parts of the province with limited survey efforts (Government of Alberta, Sustainable Resource Development 2010). Recent detections in central Alberta and the lower Athabasca Region bring the known range further east and north than previously thought, but it is unclear whether those new records represent range expansion or are simply a reflection of increased survey effort associated with resource development (C. Paszkowski, pers. comm. 2012 in COSEWIC 2012).

The population trends of Western Toads at the northern limits of their distribution in Yukon, Northwest Territories, and extreme northern British Columbia are unknown. At a monitoring site in northwestern British Columbia, the population appears to have declined with no breeding observed for three consecutive years (from 2007-2009; Slough 2009b).

3.3. Needs of the Western Toad

3.3.1. Biological Needs

The Western Toad occurs in a wide range of habitats (COSEWIC 2002) and has a biphasic life cycle consisting of aquatic eggs and tadpoles and terrestrial juveniles and adults. It requires suitable water bodies for breeding, and terrestrial habitats for foraging and hibernation as well as connectivity among seasonally used habitats (COSEWIC 2012).

Spring/breeding:

In spring, the Western Toad breeds in a variety of temporary and permanent ponds and shallow littoral zones of lakes, often with a sandy or silty bottom substrate (COSEWIC 2002; 2012). Permanent waterbodies with higher daytime temperatures (but still high dissolved oxygen concentrations) are required to promote tadpole growth (Ultsch et al. 1999; COSEWIC 2012). In the northern portion of their Canadian range, the Western Toad breeds in beaver ponds (Slough and Mennell 2006; Stevens et al. 2006), shallow stream deltas on lakes, geothermal springs, and small water bodies in gravel quarry sites (Schock 2009; Slough 2009a).

The Western Toad gathers at breeding sites to complete mating and egg-laying during a 1-2 week period in the spring. In south-central British Columbia, this is typically in late April to May and coincides with the daily minimum and maximum temperatures reaching 0 °C and 10 °C, respectively (Gyug 1999). Conversely, further north in the boreal and parkland regions of Alberta, breeding typically occurs from mid-May to early June (Browne pers. comm, 2019). The Western Toad shows some fidelity to traditional

breeding sites (Bull and Carey 2008), which may result in the selection of only one or a few of the potential breeding sites available in large areas (Slough 2004). The female lays a clutch of thousands of eggs, often communally with other females. Tadpoles often form large, dense aggregations in shallow, warm water. They develop rapidly and metamorphose in the summer within three months of egg-laying (Stebbins 1951).

Foraging/hibernation:

In the summer, adults and juveniles forage in forests and forest openings, shrubby areas, marshes, and other open and wooded habitats, and may be found several kilometers away from water bodies (COSEWIC 2002). Browne and Paszkowski (2014) found Western Toads more often in open habitat (especially wet shrubland) than predicted by habitat availability, and this was most evident during the foraging season.

Characteristics of hibernation habitats are not well known. In Alberta, hibernation sites were located in natural habitats, especially in coniferous forest stands, as opposed to human-modified or open habitats (Browne 2010). Other recent research found Western Toads hibernating in pre-existing cavities, such as rodent burrows, squirrel middens, under fallen logs of varying decay, in rocky talus slopes, or other subterranean crevices (Browne and Paszkowski 2010a; Dulisse et al. 2016). Sites must be below the frost line, as the Western Toad is intolerant of freezing, and must contain sufficient moisture to prevent desiccation (COSEWIC 2012). Most northern records come from valleys that receive consistently early, high snowfall accumulations (Cook 1977; Mennell 1997). In one Alberta study, 68% of Western Toads hibernated communally (Browne and Paszkowski 2010b).

Movement:

Newly-transformed Western Toads form large, dense, post-metamorphic aggregations along shorelines and migrate *en masse* towards terrestrial foraging areas. Terrestrial movements of the Western Toad vary with the configuration and quality of seasonal habitats. On Vancouver Island (Davis 2000) and in east-central Alberta (Browne 2010), most Western Toads used terrestrial habitats within 2 km of breeding sites although much longer movements occasionally have been reported. In Alberta, movements to reach upland hibernation sites were greater than those associated with foraging areas (Browne 2010). Hibernation sites have been found 146-1936 m (Browne and Paszkowski 2010b) and 180-6230 m (Bull 2006) from breeding sites in Alberta and Oregon, respectively. In south-central British Columbia, Dulisse et al. (2016) observed Western Toad hibernation sites ranging between 23-716 m from the lake where breeding occurs.

The Western Toad is thought to be distributed as a series of relatively independent subpopulations within a region (Davis 2002), but few data are available on dispersal, movements among breeding sites, and population dynamics.

3.3.2. Limiting Factors

Extensive movements in terrestrial habitats and long distances between breeding, hibernation, and foraging areas (Davis 2002) increase the vulnerability of Western Toads in fragmented habitats (COSEWIC 2002). Other biological factors that increase their vulnerability include communal hibernation and egg-laying (Provincial Western Toad Working Group 2014), high breeding site fidelity (Davis 2000), and the formation of large post-metamorphic aggregations (Livo 1998). Wide fluctuations in breeding success between years also increase the vulnerability of populations to local extirpation (Marsh and Trenham 2001). The species' reliance on high adult survival to sustain populations through periods of poor reproductive success means that threats that impact adult survival can have particularly pronounced effects on the population (COSEWIC 2002).

4. Threats

Threats are defined as the proximate activities or processes that have caused, are causing, or may cause in the future the destruction, degradation, and/or impairment of the entity being assessed (population, species, community, or ecosystem) in the area of interest (global, national, or subnational) (adapted from Salafsky et al. 2008). For purposes of the threat assessment (Table 2 and 3), only present and future threats are considered⁴. Threats presented here do not include limiting factors⁵, which are presented in Section 3.3.2.

⁴ Past threats may be recorded but are not used in the calculation of threat impact. Effects of past threats (if not continuing) are taken into consideration when determining long-term and/or short-term trend factors (Master et al. 2012).

⁵ It is important to distinguish between limiting factors and threats. Limiting factors are generally not human induced and include characteristics that make the species or ecosystem less likely to respond to recovery/conservation efforts (e.g., inbreeding depression, small population size, and genetic isolation; or likelihood of regeneration or recolonization for ecosystems).

4.1. Threat Assessment

The Western Toad threat assessment is based on the IUCN-CMP (World Conservation Union–Conservation Measures Partnership) unified threats classification system. Threats are defined as the proximate activities or processes that are causing, or may cause future destruction, degradation, and/or impairment of the entity being assessed (population, species, community, or ecosystem) in the area of interest (global, national, or subnational). Limiting factors are not considered during this assessment process. Historical threats, indirect, or cumulative effects of the threats, or any other relevant information that would help understand the nature of the threats are presented in the Description of Threats section. Threats for the Western Toad were assessed for the entire Canadian range (Non-calling population: Table 2 and Calling population: Table 3).

Table 2. Threat assessment table for the Western Toad Non-calling population.

Threat	Impact ^a	Scope ^b	Severity ^c	Timing ^d	
1	Residential & commercial development	Negligible	Negligible	Extreme	High
1.1	Housing & urban areas	Negligible	Negligible	Extreme	High
1.2	Commercial & industrial areas	Negligible	Negligible	Extreme	High
1.3	Tourism & recreation areas	Negligible	Negligible	Moderate	High
2	Agriculture & aquaculture	Negligible	Large	Negligible	High
2.1	Annual & perennial non-timber crops	Negligible	Negligible	Serious	High
3	Energy production & mining	Negligible	Negligible	Serious	High
3.1	Oil & gas drilling	Negligible	Negligible	Serious	High
3.2	Mining & quarrying	Negligible	Negligible	Serious	High
3.3	Renewable energy	Negligible	Negligible	Slight	High
4	Transportation & service corridors	Medium - Low	Restricted	Extreme-Moderate	High
4.1	Roads & railroads	Medium - Low	Restricted	Extreme-Moderate	High
4.2	Utility & service lines	Negligible	Negligible	Slight	High
5	Biological resource use	Low	Small	Moderate	High
5.3	Logging & wood harvesting	Low	Small	Moderate	High
6	Human intrusions & disturbance	Negligible	Negligible	Slight	High
6.1	Recreational activities	Negligible	Negligible	Slight	High
7	Natural system modifications	Negligible	Negligible	Moderate	High
7.1	Fire & fire suppression	Negligible	Negligible	Moderate	High
7.2	Dams & water management/use	Negligible	Negligible	Slight	High
7.3	Other ecosystem modifications	Negligible	Negligible	Slight	High

8	Invasive & other problematic species & genes	Medium - Low	Pervasive	Moderate - Slight	High
8.1	Invasive non-native/alien species	Medium - Low	Pervasive	Moderate - Slight	High
9	Pollution	Low	Small	Slight	High
9.1	Household sewage & urban water waste	Negligible	Negligible	Slight	High
9.2	Industrial & military effluents	Negligible	Negligible	Slight	High
9.3	Agricultural & forestry effluents	Negligible	Negligible	Slight	High
9.4	Garbage & solid waste	Negligible	Negligible	Slight	High
9.5	Air-borne pollutants	Low	Small	Slight	High
11	Climate change & severe weather	Low	Small	Moderate	High
11.2	Droughts	Low	Small	Moderate	High

Table 3. Threat assessment table for the Western Toad Calling population.

Threat		Impact ^a	Scope ^b	Severity ^c	Timing ^d
1	Residential & commercial development	Negligible	Negligible	Extreme	High
1.1	Housing & urban areas	Negligible	Negligible	Extreme	High
1.2	Commercial & industrial areas	Negligible	Negligible	Extreme	High
1.3	Tourism & recreation areas	Negligible	Negligible	Moderate	High
2	Agriculture & aquaculture	Low	Restricted	Serious - Moderate	High
2.1	Annual & perennial non-timber crops	Low	Restricted	Serious - Moderate	High
3	Energy production & mining	Low	Restricted	Serious	High
3.1	Oil & gas drilling	Low	Restricted	Serious	High
3.2	Mining & quarrying	Negligible	Negligible - Restricted	Serious	High
3.3	Renewable energy	Negligible	Negligible	Slight	High
4	Transportation & service corridors	Medium - Low	Restricted - Small	Extreme - Moderate	High
4.1	Roads & railroads	Medium - Low	Restricted - Small	Extreme - Moderate	High
4.2	Utility & service lines	Low	Small	Slight	High
5	Biological resource use	Low	Small	Moderate	High
5.3	Logging & wood harvesting	Low	Small	Moderate	High
6	Human intrusions & disturbance	Negligible	Negligible	Slight	High
6.1	Recreational activities	Negligible	Negligible	Slight	High

7	Natural system modifications	Negligible	Negligible	Moderate	High
7.1	Fire & fire suppression	Negligible	Negligible	Moderate	High
7.2	Dams & water management/use	Negligible	Negligible	Slight	High
7.3	Other ecosystem modifications	Negligible	Negligible	Slight	High
8	Invasive & other problematic species & genes	Medium - Low	Pervasive	Moderate -Slight	High
8.1	Invasive non-native/alien species	Medium - Low	Pervasive	Moderate -Slight	High
9	Pollution	Low	Restricted - Small	Moderate - Slight	High
9.1	Household sewage & urban water waste	Negligible	Negligible	Slight	High
9.2	Industrial & military effluents	Low	Small	Slight	High
9.3	Agricultural & forestry effluents	Low	Small	Slight	High
9.4	Garbage & solid waste	Negligible	Negligible	Slight	High
9.5	Air-borne pollutants	Low	Small	Slight	High
11	Climate change & severe weather	Low	Small	Moderate	High
11.2	Droughts	Low	Small	Moderate	High

^a **Impact** - The degree to which a species is observed, inferred, or suspected to be directly or indirectly threatened in the area of interest. The impact of each threat is based on Severity and Scope rating and considers only present and future threats. Threat impact reflects a reduction of a species population or decline/degradation of the area of an ecosystem. The median rate of population reduction or area decline for each combination of scope and severity corresponds to the following classes of threat impact: Very High (75% declines), High (40%), Medium (15%), and Low (3%). Unknown: used when impact cannot be determined (e.g., if values for either scope or severity are unknown); Not Calculated: impact not calculated as threat is outside the assessment timeframe (e.g., timing is insignificant/negligible or low as threat is only considered to be in the past); Negligible: when scope or severity is negligible; Not a Threat: when severity is scored as neutral or potential benefit.

^b **Scope** - Proportion of the species that can reasonably be expected to be affected by the threat within 10 years. Usually measured as a proportion of the species' population in the area of interest. (Pervasive = 71-100%; Large = 31-70%; Restricted = 11-30%; Small = 1-10%; Negligible < 1%).

^c **Severity** - Within the scope, the level of damage to the species from the threat that can reasonably be expected to be affected by the threat within an 18-year or three-generation timeframe. For this species a generation time of 6 years (COSEWIC 2012) was used resulting in severity being scored over an 18-year timeframe. Usually measured as the degree of reduction of the species' population. (Extreme = 71-100%; Serious = 31-70%; Moderate = 11-30%; Slight = 1-10%; Negligible < 1%; Neutral or Potential Benefit ≥ 0%).

^d **Timing** - High = continuing; Moderate = only in the future (could happen in the short term [< 18 years or 3 generations]) or now suspended (could come back in the short term); Low = only in the future (could happen in the long term) or now suspended (could come back in the long term); Insignificant/Negligible = only in the past and unlikely to return, or no direct effect but limiting.

4.2. Description of Threats

The overall threat impact for this species is Medium-High and High for the Non-calling and Calling populations, respectively. This overall impact considers the cumulative impacts of all threats identified for each population (IUCN-CMP 2006). Transportation and service corridors and invasive and other problematic species and genes are the most significant threats for both populations. Energy production and mining, and agriculture and aquaculture are additional threats for the Calling population. Details are discussed below for threats that are identified as Low impact or greater.

IUCN-CMP Threat 2. Agriculture & Aquaculture (Low Impact: Calling Population)

Threat 2.1 – Annual & perennial non-timber crops

The conversion of wetlands to agricultural land or intensification of agricultural practices can have serious effects on the Western Toad through loss or temporary destruction of wetland and upland foraging/dispersal/hibernation habitat (COSEWIC 2002) and mortality of dispersing/foraging toads caused by machinery. Irrigation also may reduce the suitability of aquatic breeding habitats by reducing water levels in wetlands (due to ground water diversions/water table draw down) or increasing their salinity (Eaves 2004). Agricultural lands are expanding in the Aspen Parklands in central Alberta (World Wildlife Fund 2014). Although the Western Toad may use some types of cultivated areas for foraging, these uniform habitats are unlikely to provide for seasonal needs of the Western Toad, including hibernation sites (Browne 2010).

IUCN-CMP Threat 3. Energy Production & Mining (Low Impact: Calling Population)

Threat 3.1 – Oil & gas drilling

Oil and gas exploration and development disturbs habitat of the Western Toad via ecosystem conversion, fragmentation, and environmental contamination (for impacts of associated roads and transportation infrastructure see IUCN-CMP threat #4). The oil and gas industry is most active in northeast British Columbia and north central to northern Alberta and overlaps portions of the Western Toad's range, particularly the Calling population (Austin et al. 2008). However, the footprint is relatively small relative to the range of the species, so the overall impact is predicted to be low.

IUCN-CMP Threat 4. Transportation & Service Corridors (Medium - Low Impact: Non-calling and Calling Populations)

Threat 4.1 – Roads & railroads

Roads and other transportation corridors are a threat, particularly those close to breeding sites in developed areas.. Not only do vehicles kill Western Toads crossing roads, but road construction results in a loss of woodland habitat and creates barriers to dispersal. Roads alter wetland hydrology causing hydroperiods to be out of sync with

the needs of amphibians. A negative association between abundance of aquatic-breeding amphibians and traffic volumes can be attributed primarily to road kill (Fahrig et al. 1995; Eigenbrod et al. 2008). The Western Toad often moves relatively long distances between foraging, hibernating, and breeding sites, and, hence, is more susceptible to road mortality than more sedentary amphibians (Carr and Fahrig 2001). These terrestrial movements are typically a nocturnal behaviour, therefore Western Toads are often most susceptible to road mortality at night (Dulisse et al. 2016). Although many juvenile Western Toads are vulnerable to road mortality during mass migration from natal sites (Carr and Fahrig 2001; Dulisse et al. 2016), mortality of mature females may likely have a larger impact on the population (Govindarajulu, pers. comm. 2015). Adverse effects are most severe on well-used highways, but problem areas may exist on networks of logging activities, oil and gas development, and mining roads that are expanding with increased resource extraction in many parts of the species' range in British Columbia, Alberta, and Northwest Territories. Furthermore, mortality from vehicles is not restricted to roads; for example, in the Northwest Territories, all-terrain-vehicle (ATV) tracks have been observed intersecting breeding ponds (Gau, pers. comm. 2011). Off-road vehicle use is a rapidly growing recreational sector throughout much of the Western Toad's range in Canada.

Beaver control is used to prevent road flooding and can result in loss and deterioration of beaver ponds, which are used by Western Toad as breeding habitat (Stevens et al. 2007). Irrigation ponds and borrow pits created during road construction act as population sinks for Western Toad because they provide poor larval habitat (Stevens et al. 2006).

Threat 4.2 – Utility & service lines

Seismic lines and pipelines associated with the oil and gas industry are prevalent in parts of the Western Toad range, particularly in northeastern British Columbia and north central and northern Alberta (primarily affecting the Calling population). Electrical transmission lines also intersect suitable habitat throughout the species' range. New construction of utility/service lines/corridors will impact Western Toad through habitat loss, and ongoing operation/maintenance of existing infrastructure (e.g., routine vegetation maintenance along rights-of-way) can disturb or harm dispersing/foraging individuals. However, the footprint is relatively small relative to the range of the species (e.g., the energy sector, including pipelines, seismic lines, and transmissions lines, represent 1.9% of the area of Alberta; Alberta Biodiversity Monitoring Institute 2018), so the overall impact is predicted to be low.

IUCN-CMP Threat 5. Biological Resource Use (Low Impact: Non-calling and Calling Populations)

Threat 5.3 – Logging & wood harvesting

The Western Toad is relatively tolerant of logging, but it is unclear what the long-term effects of forest harvesting might be on population dynamics. The increased proportion

of closed canopy, young second-growth stands could decrease suitable toad habitat over the long term although toads preferentially used recent clearcuts on Vancouver Island (Davis 2000). The major impact of forest harvesting on pond breeding amphibians might be the creation of breeding ponds in clearcuts that act as population sinks due to short hydroperiods (Gyug 1999).

IUCN-CMP Threat 8. Invasive & Other Problematic Species & Genes (Medium - Low Impact: Non-calling and Calling Populations)

Threat 8.1 – Invasive non-native/alien species

Epidemic diseases, particularly chytridiomycosis caused by the fungus *Batrachochytrium dendrobatidis*, threaten the Western Toad throughout its Canadian range (COSEWIC 2012). Globally, precipitous declines and extirpations of amphibians and the decline of the Western Toad in the United States have been linked to chytridiomycosis (Carey 1993; Daszak et al. 1999; Muths et al. 2003). The fungus is widely distributed within the Western Toad's Canadian range (Richardson et al. 2014): in Alberta at 7 of 15 sites surveyed (COSEWIC 2012); in the Northwest Territories (NWT) in the Fort Liard area (Schock et al. 2010); and throughout British Columbia (Adams et al. 2007; Deguise and Richardson 2009a; Slough 2009b). There are a few reports of population declines in Canada associated with the presence of the fungus (COSEWIC 2012), but even mass mortalities can easily escape detection if populations are not specifically monitored. In Atlin (northwestern British Columbia), chytridiomycosis was present in a monitored population, and might have led to the extirpation of that population (Slough 2009b). In other studies, however, Western Toads that tested positive for *B. dendrobatidis* showed no signs of disease (Deguise and Richardson 2009a; Schock et al. 2010). It is possible that unknown cofactors need to be present for the fungi to become pathogenic (Carey 1993), or that populations stressed by habitat degradation and/or increased UV-B light exposure may be more susceptible to infection (COSEWIC 2002). As humans continue to encroach on Western Toad habitats and open up new areas for development and resource extraction, the potential for introducing chytrid fungi and other pathogens increases. Disease may be transmitted by sport fishers, researchers, and others who enter water at breeding sites and travel from site to site is of concern, if equipment is not properly cleaned (British Columbia Ministry of Environment 2008; Mendez et al. 2008; Vredenburg et al. 2010).

Ranavirus (Family: Iridoviridae) is a potential threat to the Western Toad. It has caused infection and mortality in captivity and the wild (Miller et al. 2011). The distribution of ranavirus overlaps southwestern NWT where it was detected in Wood Frogs but not in Western Toads or Boreal Chorus Frogs (Schock 2009; Schock et al. 2010). No mass mortality events directly attributed to ranavirus have been documented in British Columbia nor is there any information on the prevalence of ranavirus in Western Toads in British Columbia (Govindarajulu 2007). In the Yukon, out of 19 sites containing Western Toads, eDNA sampling did not detect the presence of either pathogen (ranavirus or chytrid fungus; Hobbs and Golberg 2016). Overall, there is less evidence

for mass mortalities via ranavirus than via *Batrachochytrium dendrobatidis* (Green et al. 2002; Daszak et al. 2003 in Govindarajulu 2007).

COSEWIC (2002; 2012) identified the introduction of sport fish to previously fishless lakes as an important threat to the Western Toad. This practice is widespread through most of the species' Canadian range. Introduced fish threaten the Western Toad primarily through potential disease transmission rather than through predation, as adult and tadpole Western Toads are largely unpalatable to fish. Fish transmit pathogens, such as the water mould (*Saprolegnia* spp.), to amphibians (Kiesecker et al. 2001), but their role in chytrid transmission is unknown.

The spread of introduced species, such as Bullfrogs (*Lithobates catesbeianus*) in southwestern British Columbia, and increased abundance of predators such as raccoons (*Procyon lotor*), black and Norway rats (*Rattus* spp.), and common ravens (*Corvus corax*) in human modified landscapes are a threat to the Western Toad in localized areas (COSEWIC 2002; 2012). Predators such as raccoons and rats have been introduced to Haida Gwaii (COSEWIC 2002) and have been documented to prey upon Western Toads (Burles et al. 2004;)

IUCN-CMP Threat 9. Pollution (Low Impact: Non-calling and Calling Populations)

Threats 9.2 – Industrial & military effluents, 9.3 – Agriculture & forestry effluents, and 9.5 – Airborne pollutants

Scant research has been undertaken on the effects of specific contaminants on the Western Toad. Amphibians are vulnerable to environmental contaminants (including pesticides, herbicides, fertilizers, and road salt) that can cause a variety of effects including loss of prey base (Relyea and Diecks 2008), immunosuppression (Fontenot et al. 1994), developmental abnormalities (Kiesecker 2002; Hayes 2004; Sanzo and Hecnar 2005), and mortality (Harfenist et al. 1989; Rouse et al. 1999; Hatch et al. 2001). Contaminants from agricultural and urban sources impact Western Toad breeding sites in the South coast region and southern interior of British Columbia (Provincial Western Toad Working Group 2014). Glyphosate herbicides used for conifer release in British Columbia produce lethal and sublethal effects in amphibians (Govindarajulu 2008). Western Toads are at risk of exposure to heavy metals in tailings ponds (Brinkman 1998) and through aerial deposition (COSEWIC 2012). Exposure to heavy metals (including zinc, cadmium, and copper) increases mortality and alters growth rates of amphibians, which indirectly reduces survival (Glooschenko et al. 1992; Brinkman 1998; Bridges 2000). Acidification of wetlands from airborne sulphur associated with oil and gas extraction in northeast British Columbia and Alberta can disrupt development (Austin et al. 2008). Heavy metals and UV-B radiation may act synergistically with other environmental stressors and depress the immune system of the Western Toad; thus making them vulnerable to pathogens (Carey 1993) and deformities (Worrest and Kimeldorf 1975).

Impacts of various pollutants can be locally significant, but, over the entire range of the two populations, the overall impact is predicted to be low.

IUCN-CMP Threat 11. Climate Change & Severe Weather (Low Impact: Non-calling and Calling Populations)

Threat 11.2 – Droughts

Increased frequency and duration of droughts, predicted under climate change scenarios (IPCC 2014), can decrease the persistence of smaller wetlands, used for breeding and connectivity, and micro-sites used for rehydration (Provincial Western Toad Working Group 2011). Forest habitats may be lost as wildfires become more frequent (Guscio et al. 2008).

Climate change impacts may become more significant in the future, but, over the next 18 years (3 generations), the impact is predicted to remain low.

5. Management Objective

The management objective for the Western Toad is to maintain stable or increasing populations distributed throughout the species' present range in Canada.

Rationale for management objective

The Western Toad is widespread throughout much of western Canada. The numbers of localities and individuals are sufficiently large to maintain the viability of the species in Canada if threats are addressed and further declines are prevented. Quantifying population and habitat targets is not feasible at this time due to lack of adequate baseline information.

6. Broad Strategies and Conservation Measures

6.1 Actions Already Completed or Currently Underway

Management planning:

- A British Columbia management plan is complete (Provincial Western Toad Working Group 2014).
- Best Management Practices for amphibians and reptiles in urban and rural areas have been prepared (Ovaska et al. 2004).
- Research priorities for the Western Toad in British Columbia previously have been identified (Davis 2002).
- A Yukon amphibian management plan, including the Western Toad, is complete (Government of Yukon 2013).

- A Northwest Territories management plan for amphibians, including the Western Toad, is complete (Conference of Management Authorities 2017).

Surveys and monitoring:

Alberta:

- The Alberta Volunteer Amphibian Monitoring Program (AVAMP) was initiated in 1992 and is run by the Alberta Conservation Association (Kendall, pers. comm. 2014). In partnership with a local naturalist group, the Alberta Conservation Association runs a boreal toad monitoring project in Crowsnest Pass, southwestern Alberta (Kendall, pers. comm. 2014). All AVAMP data is compiled and submitted to Alberta Environment and Sustainable Resource Development and entered into the Fish and Wildlife Management Information System database (Kendall, pers. comm. 2014). AVAMP is developing web-based mapping products to display and disseminate volunteer-collected data from Alberta (Kendall, pers. comm. 2014).
- The Researching Amphibian Numbers in Alberta (RANA project) was operational between 1997 and 2008 and included intensive monitoring of at least four sites within the Western Toad's range (Government of Alberta, Environment and Sustainable Resource Development 2009; Wilkinson, pers. comm. 2010). Two of these sites still receive limited monitoring by volunteers and Government of Alberta staff (Wilkinson, pers. comm. 2014). Western Toads were sampled for chytrid fungus infection at 15 sites throughout their Alberta range in 2008 (Prescott, pers. comm. 2011).
- Banff National Park has extensive survey data on all amphibians and reptiles, including the Western Toad, from 1992 to 2006 (e.g., Mclvor and Mclvor 2006). In 2016, an amphibian breeding monitoring program, including Western Toads, was initiated within the park (Petersen and Rogala, pers. comm. 2019). An amphibian monitoring program in Jasper National Park, initiated in 2004, provides data on Western Toad populations in poorly known, high-elevation habitats (Shepherd, pers. comm. 2010). In Wood Buffalo National Park, inventories using automated recorders and opportunistic visits have not yielded any Western Toad records (Kindopp, pers. comm. 2011). In Waterton Lakes National Park, monitoring of sub-alpine, montane and foothills parkland has been ongoing since 1993 (Johnson 2014).
- Elk Island National Park, north of Edmonton, began instituting (spring 2016) an amphibian monitoring project. The park surrounding Beaver Hills was named a UNESCO Biosphere Reserve in 2016 (Ovaska pers. comm. 2016).

British Columbia:

- Western Toad inventories were undertaken in Mount Revelstoke and Glacier National Parks in 2003 and 2004. A long-term pond-breeding amphibian monitoring program that includes Western Toad was developed in those parks in 2009 and is ongoing (Larson and Samson 2018). Similar monitoring programs were initiated in Kootenay National Park in 2009 and Yoho National Park in 2010 (Petersen, pers. comm. 2011). Testing for presence of chytrid fungus also

occurred at a sample of Kootenay National Park sites in 2010. Since 2005, Gwaii Haanas National Park Reserve has been annually monitoring Western Toad populations at five known breeding sites (Stewart, pers. comm. 2015). During the same surveys, presence of non-native amphibians is recorded and water samples are taken for water quality.

- A draft monitoring program for Western Toads in British Columbia has been designed (Wind 2009).
- Distribution records for the province have been collated and mapped (Govindarajulu, pers. comm. 2010).
- The British Columbia Frogwatch program (British Columbia Ministry of Environment) includes a database and website for reporting occurrence records of amphibians (Surveillance for Amphibian Mass Mortalities; SAMM).
- A 5-year (2011-2015) community-based amphibian monitoring program was completed in the Nicola region in south-central BC with the Western Toad as a focal species. This work was in partnership with Biolinx Environmental Research Ltd and the Nicola Naturalist Society. Distribution within ~ 100 x 100 km study area was mapped and breeding sites were identified and monitored for multiple years (Ovaska et al. 2016).
- The ongoing (started in 2011) Summit Lake Western Toad project is located at one of the most significant known breeding sites for the species in B.C. (near Nakusp, in the southern interior of the province). Activities include highway mortality surveys and hotspot mapping, breeding site monitoring, population monitoring through mark-recapture, hibernacula research through radio-telemetry, forest harvest mitigation, camera trap monitoring of two existing highway underpasses and continual testing of various fencing materials and layout design (Dulisse et al. 2016; Dulisse pers. comm. 2019).
- Species at risk occurrence and habitat mapping (including Western Toad) was performed in the Coquitlam Watershed in 2013-2014 (Mitchell 2014).
- Amphibian inventories have been conducted and are ongoing in many parts of the province within the past 15 years (COSEWIC2012), and a study of the distribution of chytrid fungus in British Columbia amphibians has been completed (Govindarajulu et al. 2013).
- Repeat surveys at core monitoring sites were completed in 2009 and 2010 in both Mount Revelstoke and Glacier National Parks (Provincial Western Toad Working Group 2014).
- Wind and Wilmott (2012) identified known breeding sites on Vancouver Island and documented road mortality sites and mitigation efforts in B.C.
- A multi-year project investigating winter movement patterns and habitat requirements including characterization and spatial location of Western Toad hibernacula was conducted on south Vancouver Island (Wind2018).

Yukon:

- Amphibian surveys have been conducted from 1973-2010 (Slough and Mennell 2006; Slough 2009b; Cannings, pers. comm. 2010).
- Chytrid fungus surveys have been conducted in northern British Columbia and southeast Yukon (Slough 2009b). Limited monitoring of known sites is taking

place in northwestern British Columbia and southeastern Yukon, including the use of eDNA (Cannings, pers. comm. 2014; Hobbs and Goldberg 2016). These data are housed by the Yukon Conservation Data Centre, which actively comments on development proposals that may affect Western Toad habitat.

Northwest Territories:

- Amphibian population and pathogen surveys were conducted in the Sahtu and Dehcho regions in 2007 and 2008 (Schock 2009; Schock et al. 2010).
- In Nahanni National Park Reserve, inventories using automated recorders and opportunistic visits have not yielded any Western Toad records (D. Tate, pers. comm. 2011).
- The Department of Environment and Natural Resources, Government of the Northwest Territories, maintains a database of all recorded occurrences of Western Toad from the Northwest Territories.

Research:

- University of Alberta studies include projects on amphibian habitat associations (Eaves 2004; Macdonald et al. 2006; Stevens et al. 2007; Browne 2010), terrestrial movements of the Western Toad (Browne 2010), and the use of Automated Recording Units (ARUs) for amphibian monitoring (Paszkowski, pers. comm. 2014).
- Habitat use and movements have been studied in relation to resource extraction by Innovates - Technology Futures (Eaton, pers. comm. 2010).
- The ecology and movements of Western Toads have been studied on Vancouver Island (Davis 2000) and in the lower Fraser Valley (Deguise and Richardson 2009b).
- Larval growth and development in relation to sedimentation have been studied (Wood and Richardson 2009).
- The effects of mechanical forest thinning and fire on amphibians were studied at wetlands in Banff National Park, 2004-2008 (Lepitzki and Lepitzki 2003; Lepitzki and Lepitzki 2007; Maxcy and Symes 2009).
- Alberta Conservation Association and Government of Yukon are piloting the use of eDNA sampling to detect amphibian presence in freshwater wetlands (Paszkowski, pers. comm. 2014; Bennett, pers. comm. 2015).

Species and population management:

Bullfrog management activities are ongoing in the Okanagan, Kootenays and Vancouver Island (B.C.) (Govindarajulu, pers. comm. 2010 in Provincial Western Toad Working Group 2014; Fraser, pers. comm. 2019).

6.2 Broad Strategies

The following broad strategies will be used to achieve the management objective for the Western Toad:

1. Identify and secure regionally important breeding sites and terrestrial habitats.
2. Mitigate threats to important breeding sites and surrounding terrestrial habitats.
3. Undertake research to fill key knowledge gaps.
4. Establish regionally based monitoring programs to assess population status and trends.
5. Conduct outreach and stewardship.

6.3 Conservation Measures

Table 4. Conservation measures and implementation schedule for the Western Toad (Calling and Non-calling populations).

Conservation Measure	Priority ^a	Threats or concerns addressed	Timeline ^b
1. Identify and secure regionally important breeding sites and terrestrial habitats			
Using existing data and new surveys, identify and prioritize regionally important breeding sites and terrestrial habitats.	High	All	2020-2023
Secure regionally important breeding sites and terrestrial habitats, according to priority, by protecting ^c areas from the impacts of development, resource extraction and intensive human uses, and from the introduction of alien invasive species. Identify alternative breeding sites across the landscape at these priority sites for protection to ensure that metapopulation processes are preserved.	High	All	ongoing
2. Mitigate threats to important breeding sites and surrounding terrestrial habitats			
Control spread of disease and invasive species into breeding sites; establish hygiene protocols and outreach for all those who work and recreate in and around breeding sites.	High	8.1 – invasive non-native/alien species 11 – climate change	ongoing
Avoid drainage of wetlands in areas occupied by the species.	High	2.1 – annual & perennial non-timber crops 3.1 – oil & gas drilling 4.1 – roads & railroads 5.3 – logging & wood harvesting	ongoing
Develop and implement mitigation measures and Best Management Practices associated with agriculture, oil and gas/mining, logging, and sport-fishing.	Medium	2.1 – annual & perennial non-timber crops 3.1 – oil & gas drilling 4.1 – roads & railroads 5.3 – logging & wood harvesting 8.1 – invasive non-native/alien species	2020-2023
Mitigate road mortality and barriers to movements along transportation corridors.	Medium	4.1 – roads & railroads	2020-2024
3. Undertake research to fill key knowledge gaps			
Investigate factors that trigger epidemics due to chytrid fungus and other amphibian pathogens. Define current range/extent of the fungus and predicted effects of climate change.	High	8.1 – invasive non-native/alien species	2020-2028
Investigate mechanisms of spread of pathogens, including the role of sport fish introductions.	Medium	8.1 – invasive non-native/alien species	2020-2023
Investigate the impact chytrid fungus has on the age structure, and thus long-term viability, of populations that may be affected.	Medium	8.1 – invasive non-native/alien species	2020-2023
Study metapopulation dynamics, movements, and dispersal patterns in fragmented landscapes. Specifically, recording the number and range of different ponds/wetlands a population may use over the long term.	Medium	Knowledge gaps	2020-2023
Study terrestrial habitat use and identify features of hibernation sites in various landscapes; develop habitat models for mapping.	Medium	Knowledge gaps	ongoing
Clarify threats in relation to land uses, including impacts of agriculture, forestry, and energy production.	Low	Knowledge gaps	2020-2028

Conservation Measure	Priority ^a	Threats or concerns addressed	Timeline ^b
Clarify the distribution of Calling and Non-calling populations in Canada and extent of species diversity.	Low	Knowledge gaps	ongoing
Clarify the productivity and viability of constructed wetlands as means to habitat enhancement/mitigation efforts for the recovery of Western Toad populations.	Low	Knowledge gaps	ongoing
4. Establish regionally based monitoring programs to assess population status and trends			
Develop monitoring methods including analysis of existing amphibian monitoring programs.	High	Knowledge gaps	2020-2022
Initiate population monitoring in strategic locations across the species' Canadian range.	High	Knowledge gaps	2020-2021
5. Conduct outreach and stewardship			
Conduct targeted outreach (meetings, workshops, site visits and assessments) where important habitats occur on private lands or are under resource use leases on Crown lands.	Medium	2.1 – annual & perennial non-timber crops 4.1 – roads & railroads 5.3 – logging & wood harvesting 8.1 – invasive non-native/alien species 9 - pollution	2020-2022
Effectively share information on inventory and management actions; this may include brochure(s), an interactive website, etc.	Medium	2.1 – annual & perennial non-timber crops 4.1 – roads & railroads 5.3 – logging & wood harvesting 8.1 – invasive non-native/alien species 9 - pollution	2020-2022

^a **Priority** - reflects the degree to which the measure contributes directly to the conservation of the species or is an essential precursor to a measure that contributes to the conservation of the species. High priority measures are considered those most likely to have an immediate and/or direct influence on attaining the management objective for the species. Medium priority measures may have a less immediate or less direct influence on reaching the management objective, but are still important for the management of the population. Low priority conservation measures will likely have an indirect or gradual influence on reaching the management objective, but are considered important contributions to the knowledge base and/or public involvement and acceptance of the species.

^b **Timeline** - is the approximate—implementation of this management plan and is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

^c **Protection** - in this document should not be confused with the legal protection afforded to the critical habitat of threatened and endangered species under SARA. Potential mechanisms could include land zoning, securement of private lands, conservation covenants, or voluntary stewardship agreements.

7. Measuring Progress

The performance indicator presented below provides a way to define and measure progress toward achieving the management objective:

Stable or increasing Western Toad populations have been maintained throughout the species' present range in Canada.

The following targets will be used to gauge progress until there is sufficient baseline information to ensure the management objective is reached:

1. The highest priority breeding sites have been secured.
2. Best Management Practices have been developed and are actively used (e.g., appropriate hygiene measures as described in British Columbia Ministry of Environment 2008).
3. Research has been undertaken on factors that trigger epidemics of chytrid fungus (and other important amphibian pathogens) and their mechanisms of spread, and has been initiated on other threats, metapopulation dynamics, and habitat use.
4. Population monitoring programs have been initiated in strategic locations across the species' range.
5. Outreach has been implemented to target owners/managers of important habitats on private and leased lands.

8. References

- Adams, M.J., S. Galvan, D. Reinitz, R.A. Cole, S. Pyare, M. Hahr, and P. Govindarajulu. 2007. Incidence of the fungus *Batrachochytrium dendrobatidis* in amphibian populations along the northwest coast of North America. *Herpetological Review* 38: 430-431.
- Alberta Biodiversity Monitoring Institute. 2018. The status of the human footprint in Alberta – Preliminary Report 2017. Available: <https://abmi.ca/home/publications/501-550/501> [accessed February 2019].
- Austin, M.A., D.A. Buffett, D.J. Nicolson, G.G.E. Scudder, and V. Stevens (eds.). 2008. Taking nature's pulse: the status of biodiversity in British Columbia. Biodiversity BC, Victoria, B.C. 268 pp. Available: <http://www.biodiversitybc.org/> [accessed 25 February 2012].
- Beasley, B. 2011. Unpublished data. Coastal Ecologist, Association of Wetland Stewards for Clayoquot and Barkley Sounds, Ucluelet, British Columbia.
- Blaustein, A.R., J.J. Beatty, D.H. Olson, and R.M. Storm. 1995. The biology of amphibians and reptiles in old-growth forests in the Pacific Northwest. PNW-GTR-337. U.S. Department of Agriculture, Forest Service, Portland, Oregon. 98 pp.
- Bridges, C.M. 2000. Long-term effects of pesticide exposure at various life stages of the southern Leopard frog (*Rana sphenocephala*). *Archives of Environmental Contamination and Toxicology* 39: 91-96.

- Brinkman, S. 1998. Boreal toad toxicology studies. Pp. 83-114, *in* M.S. Jones, J.P. Goettl, K.L. Scherff-Norris, S. Brinkman, L.J. Livo, and A.M. Goebel (eds.). Boreal toad research progress report 1995-1997. Colorado Division of Wildlife, Fort Collins, Colorado.
- British Columbia Conservation Data Centre. 2014. Species and Ecosystems Explorer. B.C. Ministry of Environment, Victoria, B.C. Available: <http://a100.gov.bc.ca/pub/eswp/> [accessed November 2014].
- British Columbia Ministry of Environment. 2008. Standard operating procedures: Hygiene protocols for amphibian fieldwork, 2008. Victoria, British Columbia. 8 pp.
- British Columbia Ministry of Environment. 2010. Conservation framework. B.C. Ministry of Environment, Victoria, BC. Available: <https://www2.gov.bc.ca/gov/content/environment/plants-animals-ecosystems/species-ecosystems-at-risk> [Accessed November 2014].
- Browne, C. 2010. Habitat use of the Western Toad in north-central Alberta and the influence of scale. Ph.D. dissertation, Department of Biological Sciences, University of Alberta, Edmonton, Alberta.
- Browne, C.L., and C.A. Paszkowski. 2010a. Hibernation sites of Western Toads (*Anaxyrus boreas*): Characterization and management implications. *Herpetological Conservation and Biology* 5: 49-63.
- Browne, C.L., and C.A. Paszkowski. 2010b. Factors affecting the timing of movements to hibernation sites by Western Toads (*Anaxyrus boreas*). *Herpetologica* 66: 250-258.
- Browne, C.L., and C.A. Paszkowski 2014. The influence of habitat composition, season and gender on habitat selection by Western Toads (*Anaxyrus boreas*) *Herpetological Conservation and Biology* 9(2): 417-427.
- Bull, E.L. 2006. Sexual differences in the ecology and habitat selection of western toads (*Bufo boreas*) in northeastern Oregon. *Herpetological Conservation and Biology* 1: 27-38.
- Bull, E.L., and C. Carey. 2008. Breeding frequency of western toads (*Bufo boreas*) in northeastern Oregon. *Herpetological Conservation and Biology* 3: 282-288.
- Burles, D.W., A.G. Edie, and P.M. Bartier. 2004. Native land mammals and amphibian of Haida Gwaii with management implications for Gwaii Haanas National Park Reserve and Haida Heritage Site. Parks Canada Technical reports in Ecosystem Science: Report #40. Queen Charlotte City, British Columbia.
- Conference of Management Authorities. 2017. Management Plan for amphibians in the Northwest Territories. Environment and Natural Resources, Government of the Northwest Territories, Yellowknife, NT. 73 pp.

- Carey, C. 1993. Hypothesis concerning the causes of the disappearance of Boreal Toads from the mountains of Colorado. *Conservation Biology* 7: 355-362.
- Carr, L.W. and L. Fahrig. 2001. Effect of road traffic on two amphibian species of differing vagility. *Conservation Biology* 15: 1071-1078.
- Cook, F.R. 1977. Records of the Boreal Toad from the Yukon and northern British Columbia. *The Canadian Field Naturalist* 91: 185-186.
- Corkran, C., and C. Thoms. 2006. Amphibians of Oregon, Washington, and British Columbia. A Field Identification Guide - Revised and Updated. Lone Pine Publishing, Vancouver, British Columbia. 176 pp.
- COSEWIC. 2002. COSEWIC assessment and status report on the Western Toad *Bufo boreas* in Canada. Original prepared by E. Wind and L. A. Dupuis. COSEWIC. Hull, Quebec.
- COSEWIC. 2012. COSEWIC assessment and status report on the Western Toad *Anaxyrus boreas* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xiv + 71 pp. Available: www.registrelep-sararegistry.gc.ca [accessed February 2014].
- Davis, T.M. 2000. Ecology of the Western Toad (*Bufo boreas*) in forested areas on Vancouver Island. Final Report; unpublished report for Forest Renewal B.C., Ministry of Forestry, Victoria, British Columbia.
- Davis, T.M. 2002. Research priorities for the management of the Western Toad, *Bufo boreas*, in British Columbia. B.C. Ministry of Water, Land and Air Protection, Biodiversity Branch, Victoria, BC. Wildlife Working Report WR-106. 35 pp.
- Davis, T.M., and P.T. Gregory. 2003. Decline and local extinction of the Western Toad, *Bufo boreas*, on southern Vancouver Island, British Columbia, Canada. *Herpetological Review* 34: 350-352.
- Daszak, P., L. Berger, A.A. Cunningham, A.D. Hyatt, D.E. Green, and R. Speare. 1999. Emerging infectious diseases and amphibian population declines. *Emerging Infectious Diseases* 5: 735-748.
- Daszak, P., A.A. Cunningham, and A.D. Hyatt 2003. Infectious disease and amphibian population declines. *Diversity and Distributions* 9: 141-150.
- Deguisse, I., and J.S. Richardson. 2009a. Prevalence of the chytrid fungus (*Batrachochytrium dendrobatidis*) in western toads in southwestern British Columbia, Canada. *Northwestern Naturalist* 90: 35-38.
- Deguisse, I. and J.S. Richardson. 2009b. Movement behaviour of adult western toads in a fragmented, forest landscape. *Canadian Journal of Zoology* 87: 1184-1194.

- Dulisse, J., J. Boulanger, and I. Manley. 2016. Summit Lake Western Toad project 2016 field season. Prepared for Ministry of Forests, Lands Natural Resource Operations and Rural Development. 64 pp.
- Eaton, B., and M. Hiltz. 2012. Analysis of researching amphibian numbers in Alberta (RANA) Project - amphibian captures at pitfall trapping sites 1997-2008. Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 142. Edmonton, Alberta.
- Eaves, S.E. 2004. The distribution and abundance of amphibians across land-use types in Alberta's Aspen Parkland. MSc thesis. Department of Biological Sciences, University of Alberta, Edmonton, Alberta.
- Eigenbrod, F., S.J. Hecnar, and L. Fahrig. 2008. The relative effects of road traffic and forest cover on anuran populations. *Biological Conservation* 141: 35-46.
- Fahrig, L., J.H. Pedlar, S.E. Pope, P.D. Taylor, and J.F. Wegner. 1995. Effect of road traffic on amphibian density. *Biological Conservation* 73: 177-182.
- Fontenot, L.W., G.P. Noblet, and S.G. Platt. 1994. Rotenone hazards to amphibians and reptiles. *Herpetological Review* 25: 50-156.
- Frost, D.R., T. Grant, J. Faivovich, R.H. Bain, A. Haas, C.F.B. Haddad, R.O. De Sá, A. Channing, M. Wilkinson, S.C. Donnellan, C.J. Raxworthy, J.A. Campbell, B.L. Blotto, P. Moler, R.C. Drewes, R.A. Nussbaum, J.D. Lynch, D.M. Green, and W.C. Wheeler. 2006. The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1-370.
- Glooschenko, V., W.F. Weller, P.G.L. Smith, L. Alvo, and J.H.G. Archbold. 1992. Amphibian distribution with respect to pond and water chemistry near Sudbury, Ontario. *Canadian Journal of Fisheries & Aquatic Sciences* 49 (Suppl. 1): 114-121.
- Government of Alberta (Sustainable Resource Development). 2009. Alberta's amphibian monitoring programs. Available: <http://aep.alberta.ca/fish-wildlife/wildlife-management/amphibian-monitoring/researching-amphibian-numbers.aspx> [accessed March 2010].
- Government of Alberta (Environment and Sustainable Resource Development). 2010. Boreal/Foothills Sensitive Species Guidelines. Unpublished draft.
- Government of the Northwest Territories. 2018. Species at risk in the Northwest Territories, 2018. Department of the Environment and Natural Resources, Government of the Northwest Territories, Yellowknife, NT. 108 pp. Available: <https://www.nwt-species-at-risk.ca/news/guide-species-risk-nwt-2018-edition> [accessed February 2019].

- Government of Yukon. 2013. Management Plan for Yukon Amphibians. Fish and Wildlife Branch, Yukon Department of Environment, Whitehorse, Yukon. 14 pp.
- Government of Yukon. 2014. Yukon Conservation Data Centre. Available: <http://www.env.gov.yk.ca/animals-habitat/cdc.php> [accessed November 2014].
- Govindarajulu, P. 2007. Emerging infectious diseases in British Columbia Amphibians: literature review to assess risk and develop survey recommendations. Unpub. Report. B.C. Ministry of Environment, Victoria, B.C. ii + 20 pp.
- Govindarajulu, P. 2008. Literature review of impacts of glyphosate herbicide on amphibians: what risks can the silvicultural use of this herbicide pose for amphibians in B.C.? Wildlife Report No. R-28. B.C. Ministry of Environment, Victoria, B.C. v + 79 pp.
- Govindarajulu P., C. Nelson, J. LeBlanc, W. Hintz, and H. Schwantje. 2013. *Batrachochytrium dendrobatidis* surveillance in British Columbia 2008 - 2009, Canada. British Columbia Ministry of the Environment. Report ID 34795.
- Green, D.M., and R.W. Campbell. 1984. Amphibians of British Columbia. B.C. Provincial Museum Handbook 45.
- Green, D., K.A. Converse, and A.K. Schrader. 2002. Epizootiology of sixty-four amphibian morbidity and mortality events in the USA, 1996-2001. *Annals of the New York Academy of Sciences* 969: 323-339.
- Guscio, C.G., B.R. Hossack, L.A. Eby, and P.S. Corn. 2008. Post-breeding habitat use by adult boreal toads (*Bufo boreas*) after wildfire in Glacier National Park, USA. *Herpetological Conservation and Biology* 3: 55-62.
- Gyug, L.W. 1999. Timber-harvesting effects on riparian wildlife and vegetation in the Okanagan Highlands of British Columbia. B.C. Ministry of Environment, Lands and Parks Wildlife Branch, Victoria, BC. Wildl. Bull. No. B-97. 112 pp.
- Harfenist, A., T. Power, K.L. Clark, and D.B. Peakall. 1989. A review and evaluation of the amphibian toxicological literature. Canadian Wildlife Service Technical Report Series 61.
- Hatch, A.C., L.K. Belden, E. Scheellele, and A.R. Blaustein. 2001. Juvenile amphibians do not avoid potentially lethal levels of urea on soil substrate. *Environmental Toxicology and Chemistry* 20: 2328-2335.
- Hayes, T.B. 2004. There is no denying this: defusing the confusion about atrazine. *BioScience* 54:1138-1149.
- Hobbs, J., and C. Goldberg. 2016. Western Toad and Columbia Spotted Frog eDNA inventory and method assessment. Internal report prepared for the Yukon Conservation Data Centre, Whitehorse, Yukon. 26 pp.

- IPCC (Intergovernmental Panel on Climate Change). 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri, and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland. 151 pp.
- IUCN (International Union for the Conservation of Nature). 2014. The IUCN red list of threatened species. Available: <https://www.iucnredlist.org/search/external> [accessed February 2014].
- IUCN-CMP (International Union for Conservation of Nature and Conservation Measures Partnership). 2006. IUCN - CMP unified classification of direct threats, ver. 1.0 - June 2006. Gland, Switzerland. 17 pp. Available: <http://www.conservationmeasures.org/initiatives/threats-actions-taxonomies/threats-taxonomy> [accessed April 2013].
- Johnston, B., and C. Price. 2015. Amphibian monitoring in Waterton Lakes National Park, 1997-2015. Parks Canada unpublished report. 24 pp.
- Jones, L.C., W.P. Leonard, and D.H. Olson (editors). 2005. Amphibians of the Pacific Northwest. Seattle Audubon Society, Seattle, Washington, USA. 227 pp.
- Kiesecker, J.M. 2002. Synergism between trematode infection and pesticide exposure: a link to amphibian limb deformities in nature? Proceedings of Natural Academy of Sciences 99: 9900-9904.
- Kiesecker, J.M., A.R. Blaustein, and C.L. Miller. 2001. Transfer of a pathogen from fish to amphibians. Conservation Biology 15: 1064-1070.
- Larson, L. and C. Samson. 2018. Freshwater Amphibian Occupancy Measure Assessment. Glacier National Park State of the Park Report Technical Summary. 5 pp.
- Lepitzki, D.A.W., and B. Lepitzki. 2003. Experimental design for amphibian monitoring of the Carrot Creek forest treatment area, Banff National Park. Parks Canada, Banff, Alberta. 23 pp.
- Lepitzki, D.A.W., and B. Lepitzki. 2007. FESS (Fairholme Range Environmentally Sensitive Site) Amphibian Project: Draft Progress Report, 2006. Parks Canada, Banff, Alberta. 105 pp.
- Livo, L.J. 1998. Investigation of boreal toad tadpole ecology. Pp. 115-146, in M.S. Jones, J.P. Goettl, K.L. Scherff-Norris, S. Brinkman, L.J. Livo and A.M. Goebel (eds.). Boreal toad research progress report 1995-1997. Colorado Division of Wildlife, Fort Collins, Colorado.

- Macdonald, S.E., B. Eaton, C.S. Machtans, C. Paszkowski, S. Hannon, and S. Boutin. 2006. Is forest close to lakes ecologically unique?: Analysis of vegetation, small mammals, amphibians, and songbirds. *Forest Ecology and Management* 223: 1-17.
- Marsh, D.M., and P.C. Trenham. 2001. Metapopulation dynamics and amphibian conservation. *Conservation Biology* 15: 40-49.
- Master, L.L., D. Faber-Langendoen, R. Bittman, G.A. Hammerson, B. Heidel, L. Ramsay, K. Snow, A. Teucher, and A. Tomaino. 2012. NatureServe conservation status assessments: Factors evaluating species and ecosystem risk. NatureServe, Arlington, Virginia. 64pp.
- Matsuda, B.M., D.M. Green, and P.T. Gregory. 2006. Amphibians and Reptiles of British Columbia. *Royal British Columbia Museum Handbook*, Victoria, British Columbia. 266 pp.
- Maxcy, K., and S. Symes. 2009. Fairholme ecologically sensitive site amphibian research program - final progress report 2004-2008. Prepared for Parks Canada, Banff, Alberta. 54 pp.
- Mclvor, D., and M. Mclvor. 2006. Amphibian survey in Banff National Park and adjacent areas, 2005. *Bow Valley Naturalists*, Banff, Alberta. 43 pp.
- Mendez, D., R. Webb, L. Berger, and R. Speare. 2008. Survival of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* on bare hands and gloves: hygiene implications for amphibian handling. *Diseases of Aquatic Organisms* 82: 97-104.
- Mennell, R.L. 1997. Amphibians in southwestern Yukon and northwestern British Columbia. In: *Amphibians in decline: Canadian studies of a global problem*. Edited by D. Green. *Herpetological Conservation* 1: 107-109.
- Miller, D., M. Gray, and A. Storfer. 2011. Ecopathology of ranaviruses infecting amphibians. *Viruses* 2011: 2351-2373.
- Mitchell, A.M. 2014. Species at risk conservation stewardship in the Coquitlam Watershed. Prepared for BC Hydro Fish and Wildlife Compensation Program – Coastal, Burnaby, BC. 24 pp.
- Muths, E., P.S. Corn, A.P. Pessier, and D.E. Green. 2003. Evidence for disease-related amphibian decline in Colorado. *Biological Conservation* 110: 357-365.
- NatureServe. 2014. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available: <http://www.natureserve.org/explorer> [accessed February 2014].

- Ohanjanian, P., D. Adama, and A. Davidson 2006. An amphibian inventory of the East Kootenays with an emphasis on *Bufo boreas*, 2005. Prepared for Columbia Basin Fish and Wildlife Compensation Program, Nelson, British Columbia. iii + 31 pp.
- Ovaska, K, S. Lennart, C. Engelstoft, L. Matthias, E. Wind, and J. MacGarvie. 2004. Best management practices for amphibians and reptiles in urban and rural environments in British Columbia. Ministry of Water Land and Air Protection (now Ministry of Environment), Nanaimo, British Columbia. 151 pp.
- Ovaska, K., S. Lennart, and C. Engelstoft. 2016. Community-based amphibian monitoring program in multi-use landscapes in south-central BC, 2011-2015 final report. 57 pp.
- Pauly, G.B. 2008. Phylogenetic systematics, historical biogeography, and the evolution of vocalizations in Nearctic Toads (*Bufo*). Ph.D. Dissertation, University of Texas, Austin, Texas. Xvi + 165 pp.
- Provincial Western Toad Working Group. 2014. Management Plan for the Western Toad (*Anaxyrus boreas*) in British Columbia. Prepared for the B.C. Ministry of Environment, Victoria, B.C. 29 pp.
- Relyea, R.A., and N. Diecks. 2008. An unforeseen chain of events: Lethal effects of pesticides on frogs at sublethal concentrations. *Ecological Applications* 18: 1728-1742.
- Richardson, J.M.L., P. Govindarajulu, and B.R. Anholt. 2014. Distribution of the disease pathogen *Batrachochytrium dendrobatidis* in non-epidemic amphibian communities of western Canada. *Ecography* 37: 883–893.
- Rouse, J.D., C.A. Bishop, and J. Struger. 1999. Nitrogen pollution: an assessment of its threats to amphibian survival. *Environmental Health Perspectives* 107: 799-803.
- Russell, A.P., and A.M. Bauer. 2000. The Amphibians and Reptiles of Alberta. A Field Guide and Primer of Boreal Herpetology. University of Calgary Press, Calgary, Alberta. 278 pp.
- Salafsky, N., D. Salzer, A.J. Stattersfield, C. Hilton-Taylor, R. Neugarten, S.H.M. Butchart, B. Collen, N. Cox, L.L. Master, S. O'Connor, and D. Wilkie. 2008. A standard lexicon for biodiversity conservation: unified classification of threats and actions. *Conservation Biology* 22: 897-911.
- Sanzo, D., and S.J. Hecnar. 2005. Effects of road de-icing salt (NaCl) on larval wood frogs (*Rana sylvatica*). *Environmental Pollution* 140: 247-256.
- Schank, C.M. 2008. Assessing the effects of trout stocking on native amphibian communities in small boreal foothill lakes of Alberta. M.Sc. Thesis, University of Alberta, Edmonton, Alberta. 112 pp.

- Schock, D.M. 2009. Amphibian population and pathogen surveys in the Dehcho and Sahtu, Northwest Territories, 2007 and 2008. Department of Environment and Natural Resources Government of the Northwest Territories Manuscript Report No. 206. Yellowknife, Northwest Territories.
- Schock, D.M., G.R. Ruthig, J.P. Collins, S.J. Kutz, S. Carrière, R.J. Gau, A.M. Veitch, N.C. Larter, D.P. Tate, G. Guthrie, D.G. Allaire, and R.A. Popko. 2009. Amphibian chytrid fungus and ranaviruses in the Northwest Territories, Canada. *Diseases of Aquatic Organisms* 92: 231-240.
- Schock, D.M., G.R. Ruthig, J.P. Collins, S.J. Kutz, S. Carrière, R.J. Gau, A.M. Veitch, N.C. Larter, D.P. Tate, G. Guthrie, D.G. Allaire, and R.A. Popko. 2010. Amphibian chytrid fungus and ranaviruses in the Northwest Territories, Canada. *Diseases of Aquatic Organisms* 92: 231-240.
- Slough, B.G. 2004. Western Toad inventory in the Chilkoot Trail National Historic Site, July-August 2004. Parks Canada Species at Risk Inventory Fund Project SARINV04-30. 54 pp.
- Slough, B.G. 2005. Western toad, *Bufo boreas*, stewardship in the Yukon. NatureServe Yukon, Whitehorse, Yukon. 26 pp.
- Slough, B.G. 2009a. Status report on amphibians in the Yukon (Western toad *Anaxyrus boreas*, boreal chorus frog *Pseudacris maculata*, Columbia spotted frog *Rana luteiventris*, and wood frog *Lithobates sylvaticus*). Report prepared for Fish and Wildlife Branch, Environment Yukon, Whitehorse, Yukon. Draft.
- Slough, B.G. 2009b. Amphibian chytrid fungus in Western Toads (*Bufo boreas*) in British Columbia and Yukon, Canada. *Herpetological Review* 40: 319-321.
- Slough, B.G., and R.L. Mennell. 2006. Diversity and ranges of amphibians of the Yukon Territory. *Canadian Field Naturalist* 120: 87-92.
- Species at Risk Committee. 2014. Species Status Report for Western Toad (*Anaxyrus boreas*) in the Northwest Territories. Species at Risk Committee, Yellowknife, NT. 54 pp..
- Stebbins, R.C. 1951. Amphibians of western North America. University of California Press, Berkeley, California.
- Stevens, C.E., C.A. Paszkowski and D. Stringer. 2006. Occurrence of the Western Toad and its use of 'borrow pits' in west-central Alberta. *Northwestern Naturalist* 87: 107-117.
- Stevens, C.E., C.A. Paszkowski, and A.L. Foote. 2007. Beaver (*Castor canadensis*) as a surrogate species for conserving anuran amphibians on boreal streams in Alberta, Canada. *Biological Conservation* 134: 1-13.

- Tayless, E. 2011. Western Toad population summary 2005-2011 (May 31), Lost Lake population, Whistler, B.C. Resort Municipality of Whistler, British Columbia. 6 pp.
- Taylor, M., and C.M. Smith. 2003. Northern leopard frog and western toad inventory in Waterton Lakes National Park, Alberta, in 2003. Unpublished Technical Report. Parks Canada, Waterton Park, Alberta. 82 pp.
- Ultsch, G.R., D.F. Bradford, and J. Freda. 1999. Physiology: Coping with the environment. Pp. 189-214 *in* R.W. McDiarmid and R. Altig (eds). Tadpoles: The biology of anuran larvae. The University of Chicago Press, Chicago, Illinois.
- Vredenburg, V.T., R.A. Knapp, T.S. Tunstall, and C.J. Briggs. 2010. Dynamics of an emerging disease drive large-scale amphibian population extinctions. *Proceedings of the National Academy of Sciences of the U.S.A.* 107: 9689-9694.
- Wallis, C., C. Wershler, and R. Riddell. 2002. Ecological land classification of Waterton Lakes National Park, Alberta. Vol. II: wildlife resources. Parks Canada, Waterton Park, Alberta. 258 pp.
- Wind, E. 2009. Western Toad Monitoring Program in British Columbia. Draft 2. Unpublished report prepared for the B.C. Ministry of Environment, Victoria, British Columbia. 49 pp.
- Wind, E. 2018. Western Toad winter habitat requirements in modified landscapes on Vancouver Island – Summary. Prepared for the Ministry of Forests, Lands, Natural Resource Operations and Rural Development, Nanaimo, BC. 67 pp.
- Wind, E., and S. Wilmott. 2012. Identification of Western Toad (*Anaxyrus boreas*) breeding sites and road mortality mitigation assessment on Vancouver Island. Prepared for Ministry of Forests, Lands and Natural Resource Operations. Nanaimo, BC. 12 pp..
- Wood, S., and J.S. Richardson. 2009. The impacts of fine-sediment loading on the growth and survival of western toads (*Bufo boreas*). *Freshwater Biology* 54: 1120-1134.
- Working Group on General Status of NWT Species. 2016. NWT Species 2016-2020 – General status ranks of wild species in the Northwest Territories. Department of Environment and Natural Resources, Government of Northwest Territories, Yellowknife, NT. 304 pp.
- World Wildlife Fund 2014. Canadian Aspen Forests and Parkland. Available: <http://www.worldwildlife.org/ecoregions/na0802> [accessed October 2014].
- Worrest, R.C., and D.J. Kimeldorf. 1975. Photoreactivation of potentially lethal, UV-induced damage to Boreal Toad (*Bufo boreas boreas*) tadpoles. *Life Sciences* 17: 1545-1550.

Yukon CDC (Conservation Data Centre). 2015. Report your sightings – Yukon animals of conservation concern, March 2015. Available: <http://www.env.gov.yk.ca/animals-habitat/documents/Amphibians2015.pdf>. [accessed January 2016].

9. Personal Communications

Bennett, Bruce. Yukon Conservation Data Centre Coordinator, Yukon Government, Whitehorse, YT. Personal communication through email to S. Dar. 2015.

Browne, Constance. Research Associate – Herpetology, New Brunswick Museum, Saint John, NB. Personal communication through email to J. Maida. 2019.

Cannings, Syd. Species at Risk Biologist, Environment and Climate Change Canada, Whitehorse, YT. Personal communication through email to H. Middleton. 2014.

Dulisse, Jakob. Biologist, Jakob Dulisse Consulting, Nelson, BC. Personal communication through email to J. Maida. 2019.

Eaton, Brian. Aquatic Ecology Research Scientist, Alberta Innovates - Technology Futures, Vegreville, AB. Personal communication through telephone to K. Ovaska. 2010.

Fraser, Khaylish. Aquatics Program Coordinator, Central Kootenay Invasive Species Society, Robson, BC. Personal communication through email to J. Maida. 2019.

Gau, Rob. Wildlife Biologist (Species at Risk), Northwest Territories Department of Environment and Natural Resources, Yellowknife, NT. Personal communication through email to S. Cannings. 2011.

Govindarajulu, Purnima. Small Mammal and Herpetofauna Specialist, B.C. Ministry of Environment, Victoria, BC. Personal communication through telephone to K. Ovaska. 2010 and H. Middleton. 2015.

Hughson, Ward. Aquatic Specialist, Jasper National Park, AB. Personal communication through email to H. Middleton. 2015.

Jung, Thomas. Senior Wildlife Biologist, Environment Yukon, Whitehorse, YT. Personal communication through email to S. Cannings. 2011.

Kendell, Kris. Senior Biologist, Alberta Conservation Association. Personal communication through email to H. Middleton. 2014.

Kindopp, Rhonda. Park Ecologist, Parks Canada, Fort Smith, NT. Personal communication through email via D. Casimir to S. Cannings. 2011.

- Mclvor, M. *In* COSEWIC. 2012. Bow Valley Naturalist, Banff, AB. Personal communication through email and photo of a calling Western Toad from Kootenay National Park to D. Lepitzki. 2012.
- Ovaska, Kristiina. Senior Ecologist, Biolinx Environmental Ltd., North Saanich, BC. Personal communication through comment-response table to J. Maida. 2019.
- Paszkowski, Cynthia.. *In* COSEWIC. 2012. Professor and Associate Chair Research, Department of Biological Sciences, University of Alberta, Edmonton, AB. Personal communication through email to B. Slough and K. Ovaska. 2012.
- Petersen, Derek. Conservation Biologist, Parks Canada, Radium, BC. Personal communication through email via D. Casimir to S. Cannings. 2011.
- Petersen, Derek. Ecological Integrity Monitoring Coordinator, Lake Louise, Yoho, Kootenay and Banff Field Units, Parks Canada, Radium, BC. Personal communication through email to J. Maida. 2019.
- Prescott, David. Senior Species at Risk Biologist, Alberta Sustainable Resource Development, Red Deer, AB. Personal communication through telephone to S. Cannings. 2011.
- Rogala, Kimo. Resource Management Officer II, Ecological Integrity Monitoring Specialist, Banff National Park, Parks Canada, Banff, AB. Personal communication to J. Maida. 2019.
- Russell, John. Retired Wildlife Biologist, Waterton, AB. Personal communication through email to B. Johnston. 2013.
- Shepherd, Brenda. Park Ecologist, Jasper National Park, Parks Canada, Jasper, AB. Personal communication through mail to K. Ovaska. 2010.
- Stevens, Scott. Species at Risk Biologist, Alberta Sustainable Resource Development, Red Deer, AB. Personal communication through email to K. Ovaska. 2010.
- Stewart, Heather. A/Ecologist Team Leader. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and Haida Heritage Site. Personal communication through email to Diane Casimir. 2015.
- Tate, Doug. Conservation Biologist, Parks Canada, Fort Simpson, NT. Personal communication through email via D. Casimir to S. Cannings. 2011.
- Wilkinson, Lisa. Non-game/Species at Risk Biologist, Alberta Sustainable Resource Development, Edson, AB. Personal communication through email to K. Ovaska. 2010.

Appendix A: Effects on the Environment and Other Species

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the [Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals](#)⁶. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or any of the [Federal Sustainable Development Strategy](#)'s⁷ goals and targets.

Conservation planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that implementation of management plans may also inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the management plan itself, but also are summarized in this statement.

Conservation and management of wetland breeding habitat across the range of the Western Toad will benefit a number of other species of conservation concern that also depend on wetland habitat for all or part of their lifecycle. Similarly, conservation and management of upland hibernation, foraging, and dispersal habitats is expected to have positive effects on co-occurring species. The potential for the plan to inadvertently lead to adverse effects on other species was considered. The SEA concluded that this plan will clearly benefit the environment and will not entail any significant adverse effects.

⁶ www.canada.ca/en/impact-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html

⁷ www.fsds-sfdd.ca/index.html#/en/goals/